arXiv:1505.01930v1 [math.AP] 8 May 2015

BOUNDARY VALUE PROBLEM FOR A PARABOLIC-HYPERBOLIC EQUATION IN A RECTANGULAR DOMAIN

Djumaklych Amanov
Institute of Mathematics at the National University of Uzbekistan
e-mail: damanov@yandex.ru

Abstract. In present paper we study a boundary value problem for a mixed parabolic-hyperbolic type equation in a rectangular domain and prove the existence of unique solution of this problem. In theory of boundary value problems for second order mixed type equations usually two conjugation conditions are in use. In this case, for mixed type equations containing hyperbolic equation in a rectangular domain for solvability of boundary value problem appears certain condition. In this paper we give three conjugation conditions. In this case mentioned condition not appears.

Keywords. Mixed type equation, boundary value problem, Fourier's method, uniqueness, existence.

MSC2010: 35M12.

1 Introduction

In domain $\Omega = \{(x,t) : 0 < x < p, -T < t < T\}$ we consider equation

$$\left.\begin{array}{l} u_{tt} - u_{xx}, t > 0 \\ u_t + u_{xx}, t < 0 \end{array}\right\} = f(x,t), \tag{1}$$

where $f(x,t)$ is given function.

We denote by $\Omega^+ = \Omega \cap (t > 0)$, $\Omega^- = \Omega \cap (t < 0)$.

Problem. Find in $\Omega^+ \cup \Omega^-$ solution $u(x,t) \in C\left(\overline{\Omega}\right) \cap C^{2,2}_{x,t}(\Omega)$ of equation (1) satisfying boundary conditions

$$u(0,t) = 0, u(p,t) = 0, -T < t < T \tag{2}$$

and conjugation conditions

$$u(x,+0) = u(x,-0), 0 < x < p, \tag{3}$$

$$u_t(x,+0) = u_t(x,-0), 0 < x < p, \tag{4}$$

$$u_{tt}(x,+0) = u_{tt}(x,-0), 0 < x < p. \tag{5}$$

Boundary value problems for mixed parabolic-hyperbolic type equations when hyperbolic part of domain is characteristic triangle was studied by many authors (see [1],[2],[3] and references therein). Boundary value problems for parabolic-hyperbolic equations in rectangular domains was studied by Sabitov K.B. in [4-6] and for the higher order generalized mixed parabolic equation with fractional derivatives in [7]. The problem (1)-(5) differs from the known works of the conjugation condition (5). If we refuse condition (5) and put $u(x,T) = 0$ then in Ω^+ we have Dirichlet type problem, the solvability of which is strongly depends on the relation of the sides of the rectangle [8]. In our case, this dependence is expressed by inequality $\cos\lambda_n T + \lambda_n \sin\lambda_n T \neq 0, \lambda_n = \frac{n\pi}{p}, n = 1, 2, ...$ the validity of which is not obvious. A similar inequality is given in [4], which proved is validity under certain restrictions on numbers T and p .

2 Uniqueness of solution of problem (1)-(5)

Theorem 1. Let conditions

$$\lim_{x\to 0} xu_x\left(x,t\right)=0, \lim_{x\to p}\left(p-x\right)u_x\left(x,t\right)=0 \tag{6}$$

is valid, then the solution of problem (1)–(5) is unique, if it exists.

Proof. Let $f\left(x,t\right)=0$ in $\overline{\Omega}$. We will show that $u\left(x,t\right)=0$ in $\overline{\Omega}$. Following [9] consider integrals

$$\alpha_n\left(t\right)=\int\limits_0^p u\left(x,t\right)X_n\left(x\right)dx,\quad t>0, \tag{7}$$

$$\beta_n\left(t\right)=\int\limits_0^p u\left(x,t\right)X\left(x\right)dx,\quad t<0, \tag{8}$$

where $X_n\left(x\right)=\sqrt{\frac{2}{p}}\sin\lambda_n x, \lambda_n=\frac{n\pi}{p}, n=1,2,...$

On the basis of (7) and (8) we introduce the functions

$$\alpha_{n,\varepsilon}\left(t\right)=\int\limits_\varepsilon^{p-\varepsilon} u\left(x,t\right)X_n\left(x\right)dx, \tag{9}$$

$$\beta_{n,\varepsilon}\left(t\right)=\int\limits_\varepsilon^{p-\varepsilon} u\left(x,t\right)X_n\left(x\right)dx, \tag{10}$$

where $\left(\varepsilon,p-\varepsilon\right)\neq\emptyset$. Differentiating (9) two times with respect to t we have

$$\alpha''_{n,\varepsilon}\left(t\right)=\int\limits_\varepsilon^{p-\varepsilon} u_{tt}\left(x,t\right)X_n\left(x\right)dx.$$

From homogeneous equation (1) we get $\alpha''_{n,\varepsilon}\left(t\right)=\int\limits_\varepsilon^{p-\varepsilon} u_{xx}\left(x,t\right)X_n\left(x\right)dx$. Integrating by parts last integral and using conditions (6), we find

$$\alpha''_n\left(t\right)+\lambda_n^2\alpha_n\left(t\right)=0. \tag{11}$$

Analogously we get

$$\beta'_n\left(t\right)-\lambda_n^2\beta_n\left(t\right)=0. \tag{12}$$

The general solutions of equations (11), (12) have the form

$$\alpha_n\left(t\right)=a_n\cos\lambda_n t+b_n\sin\lambda_n t,$$
$$\beta_n\left(t\right)=c_n e^{\lambda_n^2 t}$$

.

Using conjugation conditions (3)-(5) which pass the next form $\alpha_n\left(0\right)=\beta_n\left(0\right), \alpha'_n\left(0\right)=\beta'_n\left(0\right), \alpha''_n(0)=\beta''_n\left(0\right)$ we find $a_n=b_n=c_n=0$. Consequently $\alpha_n\left(t\right)=\beta_n\left(t\right)=0$. Then from completeness of functions $X_n\left(x\right)$ from (7) and (8) we obtain $u\left(x,t\right)\equiv 0$ in $\overline{\Omega}$. Theorem 1 is proved.

3 Existence of the solution of problem (1)-(5)

We denote

$$u(x,t)=\begin{cases} u^{+}(x,t)\,, & t>0, \\ u^{-}(x,t)\,, & t<0. \end{cases}$$

Using Fourier's method we get the solution of problem (1)-(5) in the form

$$u^{+}(x,t)=\sum_{n=1}^{\infty}X_n(x)\left[\frac{\left(1-\lambda_n^2\right)f_n(0)-f'{}_n(0)}{\lambda_n^2\left(\lambda_n^2+1\right)}cos\lambda_n t+\frac{2f_n(0)-f'{}_n(0)}{\lambda_n\left(\lambda_n^2+1\right)}\sin\lambda_n t+\right.$$

$$\left.\frac{1}{\lambda_n}\int\limits_0^t f_n(\tau)\sin\lambda_n(t-\tau)\,d\tau\right] \tag{13}$$

$$u^{-}(x,t)=\sum_{n=1}^{\infty}X_n\left[\frac{\left(1-\lambda_n^2\right)f_n(0)-f'{}_n(0)}{\lambda_n^2\left(\lambda_n^2+1\right)}e^{\lambda_n^2 t}-\int\limits_t^0 f_n(\tau)\,e^{\lambda_n^2(t-\tau)}d\tau\right], \tag{14}$$

We consider the following derivatives

$$\begin{gathered}\frac{\partial u^{+}}{\partial t}(x,t)=\sum_{n=1}^{\infty}X_n(x)\left[\frac{\left(\lambda_n^2-1\right)f_n(0)+f'{}_n(0)}{\lambda_n^2+1}\sin\lambda_n t+\right.\\ \left.+\frac{2f_n(0)-f'{}_n(0)}{\lambda_n^2+1}\cos\lambda_n t+\int\limits_0^t f_n(\tau)\cos\lambda_n(t-\tau)\,d\tau\right],\end{gathered} \tag{15}$$

$$\frac{\partial u^{-}}{\partial t}(x,t)=\sum_{n=1}^{\infty}X_n(x)\left[\frac{2f_n(0)-f'{}_n(0)}{\lambda_n^2+1}e^{\lambda_n^2 t}-\int\limits_t^0 f'{}_n(\tau)e^{\lambda_n^2(t-\tau)}d\tau\right], \tag{16}$$

$$\frac{\partial^2 u^{+}}{\partial t^2}(x,t)=\sum_{n=1}^{\infty}X_n(x)\left[\frac{2\lambda_n^2 f_n(0)+f'{}_n(0)}{\lambda_n^2+1}\cos\lambda_n t+\right.$$

$$\left.+\frac{\left(2\lambda_n^2+1\right)f'{}_n(0)-2\lambda^2{}_n f_n(0)}{\lambda_n\left(\lambda_n^2+1\right)}\sin\lambda_n t+\frac{1}{\lambda_n}\int\limits_0^t f''{}_n(\tau)\sin\lambda_n(t-\tau)\,d\tau\right], \tag{17}$$

$$\frac{\partial^2 u^{+}}{\partial x^2}(x,t)=\sum_{n=1}^{\infty}X_n(x)\left[\frac{2\lambda_n^2 f_n(0)+f'{}_n(0)}{\lambda_n^2+1}\cos\lambda_n t-f_n(t)+\right.$$

$$\left.+\frac{\left(2\lambda_n^2+1\right)f'{}_n(0)-2\lambda_n f_n(0)}{\lambda_n\left(\lambda_n^2+1\right)}\sin\lambda_n t+\frac{1}{\lambda_n}\int\limits_0^t f''{}_n(\tau)\sin\lambda_n(t-\tau)\,d\tau\right], \tag{18}$$

$$\frac{\partial^2 u^{-}}{\partial x^2}(x,t)=\sum_{n=1}^{\infty}X_n(x)\left[\frac{f'{}_n(0)-2f_n(0)}{\lambda_n^2+1}e^{\lambda_n^2 t}+f_n(t)+\int\limits_t^0 f'{}_n(\tau)\,e^{\lambda_n^2(t-\tau)}d\tau\right], \tag{19}$$

$$\frac{\partial^2 u^-}{\partial t^2}(x,t) = \sum_{n=1}^{\infty} X_n(x)\left[\frac{2\lambda_n^2 f_n(0) + f'_n(0)}{\lambda_n^2+1} e^{\lambda_n t} - \int_t^0 f''_n(\tau)\, e^{\lambda_n^2(t-\tau)} d\tau\right]. \qquad 20$$

Lemma 1. Let $f(x,t) \in C_{x,t}^{1,0}\left(\overline{\Omega}\right), f(0,t) = f(p,t) = 0, \frac{\partial f}{\partial x} \in Lip_\alpha[0,p], 0 < \alpha < 1$ uniformly with respect to t, then series

$$\sum_{n=1}^{\infty} f_n(t)\, X_n(x), \tag{21}$$

where

$$f_n(t) = \int_0^p f(x,t)\, X_n(x)\, dx \tag{22}$$

converges absolutely and uniformly in $\overline{\Omega}$.

Proof. Integrating by parts integral in (22), we find

$$f_n(t) = \frac{1}{\lambda_n} f_n^{(1,0)}(t)$$

where

$$f_n^{(1,0)}(t) = \int_0^p \frac{\partial f}{\partial x}(x,t) \sqrt{\frac{2}{p}} \cos\lambda_n x dx \tag{23}$$

according to [10] $\left|f_n^{(1,0)}(t)\right| \leq \frac{c}{n^\alpha}, c = const > 0$. Then $|f_n(t)| < \frac{cp}{\pi}\frac{1}{n^{1+\alpha}}$ and series $\sum\limits_{n=1}^{\infty}\frac{1}{n^{1+\alpha}}$ converges. Consequently, series in (21) absolutely and uniformly converges in $\overline{\Omega}$.

Lemma 1 is proved.

Lemma 2. If $f(x,t) \in C_{x,t}^{0,1}\left(\overline{\Omega}\right), \frac{\partial^2 f}{\partial t^2} \in L_2(\Omega)$, then following estimates are valid

$$\left|\int_t^0 f_n^{(k)}(\tau)\, e^{\lambda_n^2(t-\tau)} d\tau\right| \leq \frac{c_k}{\lambda_n}\left\|f_n^k\right\|_{L_2(-T,0)},$$

$$\left|\frac{1}{\lambda_n}\int_0^t f_n^{(k)}(\tau)\sin\lambda_n(t-\tau)\, d\tau\right| \leq \frac{\widetilde{c_k}}{\lambda_n}\left\|f_n^{(k)}\right\|_{L_2(0,T)}, k = 0,1,2,$$

where $c_k = const > 0, \widetilde{c_k} = const > 0$. The proof of this Lemma follows from Hölder inequality for integrals.

Lemma 3. If $f(x,t) \in C\left(\overline{\Omega}\right)$, $\frac{\partial f}{\partial x} \in L_2(\Omega)$, then series

$$\sum_{n=1}^{\infty} X_n(x)\frac{2\lambda_n^2 f_n(0)}{\lambda_n^2+1} cos\lambda_n t \tag{23}$$

absolutely and uniformly converges in $\overline{\Omega}$.

Proof. Integrating by parts integral $\int\limits_0^p f\left(x,0\right) X_n\left(x\right) dx$ we have $f_n\left(0\right) = \frac{1}{\lambda_n} f_n^{(1,0)}\left(0\right)$ and $\frac{2\lambda_n^2 \left|f_n(0)\right|}{\lambda_n^2+1} \leq 2\left|f_n\left(0\right)\right| = \frac{2}{\lambda_n}\left|f_n^{(1,0)}(0)\right|$. For series (23) the series $\sum\limits_{n=1}^{\infty} \frac{\left|f_n^{(1,0)}(0)\right|}{\lambda_n}$ is majorant. Applying Hölder's inequality for sums to last series we get

$$\sum_{n=1}^{\infty} \frac{\left|f_n^{(1,0)}\left(0\right)\right|}{\lambda_n} \leq \left(\sum_{n=1}^{\infty} \frac{1}{{\lambda_n}^2}\right)^{\frac{1}{2}} \left(\sum_{n=1}^{\infty} \left|f_n^{(1,0)}\left(0\right)\right|^2\right)^{\frac{1}{2}} \leq \frac{p}{\sqrt{6}} \left\|\frac{\partial f\left(x,0\right)}{\partial x}\right\|_{L_2(0,p)}.$$

Consequently series (23) absolute and uniformly converges in $\overline{\Omega}^{+}$.

Lemma 3 is proved.

Theorem 2. Let $f\left(x,t\right) \in C^1\left(\overline{\Omega}\right), \frac{\partial^2 f}{\partial t^2} \in L_2\left(\Omega\right), f\left(0,t\right) = f\left(p,t\right) = 0, \frac{\partial f}{\partial x} \in Lip_{\alpha}[0,p]$ uniformly with respect to t , then series (13)–(20) absolutely and uniformly converge in $\overline{\Omega^+}$ and $\overline{\Omega^-}$ respectively. Solutions (13) and (14) satisfy equation (1) in Ω^+ и Ω^- and conditions (2)–(5).

Proof. Deducting (18) from (17) we convinced that solutions (13) and (14) satisfied equation (1) in Ω^+. Adding (16) and (19) we convinced that solutions (13) and (14) satisfied equation (1) in Ω^-. From properties of functions $X_n\left(x\right)$ follows that solutions (13) and (14) satisfies conditions (2). Passing to limit in (13) and (14), (15) and (16), (17) and (20) as $t \to 0$, we convinced that solutions (13) and (14) satisfy the conditions (3)–(5).

Theorem 2 is proved.

Remark. In the domain Ω if we consider mixed elliptic - hyperbolic type equation

$$\left.\begin{array}{ll} u_{tt} - u_{xx}, & t > 0 \\ u_{tt} + u_{xx}, & t < 0 \end{array}\right\} = f(x,t)$$

with conditions (2) – (5) and $u(x,\ -T) = 0$, then in Ω^+ the Diriclet type problem not appears.

References

1. T.D. Dzhurayev, Boundary value problems for mixed and mixed-composite type equations.Tashkent, Fan, 1979.p.240.

2. T.D. Dzhurayev, A.Sopuyev and M. Mamajanov, Boundary value problems for equations of parabolic-hyperbolic type. Tashkent. Fan. 1986, p.220.

3. T.D. Dzhurayev and A.Sopuyev, To the theory of partial differential equations of fourth order. Tashkent, Fan, 2000. p.144.

4. K.B. Sabitov, Tricomi problem for a mixed parabolic-hyperbolic type equation in a rectangular domain. Mat. Zametki, 2008.v.86(2). pp.273–279. (in Russian)

5. K.B. Sabitov, L.H. Rakhmanova, Initial-boundary value problem for a mixed parabolic-hyperbolic type equation in a rectangular domain. Differenc. Uravneniya 2008. v. 44, №9, pp.1175–1181. (in Russian)

6. K.B. Sabitov, E.M.Safin. An inverse problem for a mixed parabolic-hyperbolic type equation in a rectangular domain. Izv.Vuzov.Matematika. 2010, №4, pp.55–62. (in Russian)

7. D. Amanov and J.M. Rassias. Boundary value problems for the higher order generalized mixed-parabolic equation with fractional derivatives. Contemporary Analysis and Applied Mathematics.vol.2, №.2, pp.198–211, 2014.

8. D. Amanov, A.Ashyralyev. Well-posedness of boundary value problems for partial differential equations of even order. Electronic Journal of Differential Equation. Vol.2014 (2014), №108, pp 1–8.

9. E.I.Moiseyev. About solution of a nonlocal boundary value problem by spectral method. Differenc. Uravneniya.1999. vol.35, №8. pp. 1094-1100.(in Russian)

10.N.K. Bari. Trigonometrical series.M.1961.(in Russian)